



Oral Parasites and its Associated with the Population Residency and Some Physiological Disorders among Population in Erbil City.

Hawri Hawar Mohammed Jabbary *, Narmin Rafiq Hamad**

*Medical Laboratory Technology, Erbil Health and Medical Technical College, Erbil Polytechnic University-Erbil

**Biology Dep., Science College, Salahhadin University-Erbil-Iraq

*Corresponding author email: hawri.mohammed@epu.edu.iq, 07504458017

| | | | | | |
|-----------|-----------|-----------|-----------|------------|------------|
| Received: | 29/7/2022 | Accepted: | 27/9/2022 | Published: | 31/12/2022 |
|-----------|-----------|-----------|-----------|------------|------------|

ABSTRACT

Background:

The parasitic infection may be caused by the parasites *Trichomonas tenax* and *Entamoeba gingivalis*, which were included in this investigation. This study's goal is to estimate the incidence of the parasites in patients' oral cavities with associated factors like residency, dental caries and diabetes in Erbil, Iraq.

Materials and Methods:

The current study comprised 354 patients in total. A questionnaire recorded residency (urban and Rural), dental caries and diabetes. Calculi samples were taken by use of a sterile curette and were placed in sterile collection vials, each containing 1 ml of sterile normal saline solution. Saliva samples of the patients were collected in sterile vials, then dilution was done by addition of 1 ml of sterile saline solution to each vial. Then each vial was labeled by patient's number, name and date of collection. The purpose of collection the calculi and saliva to search about the parasites in the oral cavity.

Results:

The higher infection ratio of *E. gingivalis* in the rural regions in comparison with that of urban areas (91.7% and 89% respectively), and the results discovered that the higher positive rate of *E. gingivalis* (93.3%) was observed among those with dental caries and it was encountered in (86.8%) of subjects free from dental caries, regarding diabetes, diabetes and the presence of *E. gingivalis* were shown to have a statistically significant connection (97.1%) and (35.7%) infection rates of *E. gingivalis* among diabetics and non-diabetic participants respectively. While *T. tenax* was more common in individuals with non-diabetes (12.2%) compared to the diabetics (9.5%), but no relation was found.

Conclusion:

Infection with these parasites is relatively prevalent in people with dental caries, diabetes, and the persons' environment. It appears that continuation of advices are important in regulator of parasitic infection in Kurdistan Region of Iraq.

Key words:

Gingivitis, *Trichomonax tenax*, oral disease.



الخلاصة

مقدمة:

تبين في هذا البحث أن الطفيليات التي تضمنتها الدراسة هي أميبا الفم *Entamoeba gingivalis* والمسوطة الفمية *Trichomonas tenax* ربما مسئولتان عن الإصابة بالطفيليات الفمية. الغرض من هذا الدراسة هي التحري عن الطفيليات الموجودة في تجويف الفم و علاقتها بمكان العيش (السكن)، تسوس الاسنان و داء السكري بين مراجعين في مراكز الاسنان في محافظة أربيل-العراق.

طرق العمل:

من مجموع 354 حالة التي تضمنتها هذا الدراسة. وكذلك الاستبيان سجلت السكن، تسوس الاسنان و مرض السكري. نموذج راسب الفم جمعت بواسطة استعمال curette معقم ووضعت في أنبوبة جمع معقم، كل منها يحتوي على (1مل) من محلول Normal saline المعقم. و نموذج اللعاب وضعت في أنبوبة معقمة، ومن ثم تم تخفيفها بواسطة إضافة (1مل) من N.S. المعقم. كتابة و تدوين رقم المريض، أسم المريض و تاريخ جمع النموذج على كل أنبوبة.

الاستنتاجات:

النسبة العالية للإصابة بالمسوطة الفمية لدى المراجعين الذين يستوطنون أطراف المحافظة بالمقارنة مع المدينة (91.7% و 89%) على التوالي. أظهرت علاقة أحصائية معنوية لأميبا الفم E.g. (93.3 %) في فم الاشخاص الذين لديهم تسوس الاسنان و بالعكس، و(86.8 %) حالة في فم الذين ليس لديهم أي تسوس. وبخصوص مرض السكري أحصائيا هناك علاقة قوية بين الاشخاص المصابين بمرض السكري والإصابة باميبا الفم *Entamoeba gingivalis* (97.1%) و (35.7%) بين المصابين بالسكري والذين ليس لديهم المرض على التوالي. بينما المسوطة الفموية *Trichomonas tenax* شائع جدا بين الاشخاص الذين ليس لديهم مرض السكري (12.2%) مقارنة مع الذين لديهم المرض (9.5%)، ولكن ليس هناك أي علاقة معنوية.

الكلمات المفتاحية:

التهاب اللثة، المسوطة الفمية، أمراض الفم



INTRODUCTION

Human oral cavity is colonized by specific bacteria, protozoa, and fungi [1]. The oral cavity is frequently rather stable; nonetheless, some circumstances may result in significant oral conditions like periodontitis and caries because several oral commensal microbes may result illnesses if suitable circumstances are supplied [2]. Some authors measured *Entamoeba gingivalis* as pathogenic while other considered the parasite as commensal [3]. Molecular approaches have lately proven the importance of bacteria in human health and illness [4].

Trichomonas tenax is one members of family Trichomonadidae, is a parasite that habitat in the oral cavity of human and is anaerobic. Many analyses were conducted aiming at determining the frequently seen the parasite in those with chronic periodontitis, it discovered accuracy in individuals with poor dental hygiene and chronic periodontitis, it is worldwide distributed, ranging from 4% to 53% [5]. Most *Trichomonas* are parasitic or commensal flagellates that is living in low-oxygen environments [6].

Both parasite is cosmopolitan and are most usually found in people who have poor oral hygiene with periodontal disease advanced [7]. The evidence from previous researches about oral protozoa are likewise restricted in some countries.

Materials and Methods

• Subjects

The current study included 354 individuals, were enrolled over a 3-month period, January to March 2022, we have also visited the College of Dentistry's dental care facilities and Xanzad Learning Center and also individuals from the Private Clinic in Erbil–Kurdistan region of Iraq, for founding the incidence of oral protozoa *E. gingivalis* and *T. tenax* in relation with residency, dental caries and diabetes among the participants. Information questionnaire form was collected from each patient.

The patients' fully informed permission was obtained. The people were chosen and tested to ensure that they met the requirements for dental caries and diabetes illness. Wet preparations are typically made from mouth scrapings [8]. Then direct examination for each slide was done by use of 400X magnification of light microscope for detection of the oral protozoans. The movement and morphological structure of the parasites were used for purpose of identify [9].



Results and Discussion

Residency

Statistically, insignificant relationship was present between both oral protozoa and residency and the findings of this study are shown in the Tables (1 and 2) respectively considering the greater infection incidence of *E. gingivalis* in rural locations in compared with that of urban regions (91.7% and 89% respectively). This can be explained by the poor socioeconomic position of the patients in these circumstances, poor hygiene, and a lack of access to public health facilities.

These findings are consistent with the findings of [10] JiaYan et al. in Tangshan, China, who stated that gingival amoebic infection was not associated with urban/rural residency. On the contrary, regarding *T. tenax*, the lower infection rate was among rural population in comparison with that of urban inhabitants (6.5% and 11.4% respectively). The current study's findings might be attributed to the involved low condition of oral hygiene, which increased the incidence of oral protozoa [11, 12, 13].

Table (1) *Entamoeba gingivalis* prevalence based on the patients' residence

| Residency | <i>Entamoeba gingivalis</i> | | |
|-----------|-----------------------------|-------------|-------------|
| | No. examined | No. +ve (%) | No. -ve (%) |
| Urban | 246 | 219 (89) | 27 (11) |
| Rural | 108 | 99 (91.7) | 9 (8.3) |
| Total | 354 | 318 (89.8) | 89.8 |

P value= 0.291

Table (2) Prevalence of *Trichomonas tenax* according to the residency of the patients

| Residency | <i>Trichomonas tenax</i> | | |
|-----------|--------------------------|-----------|------------|
| | No. examined | No. +ve | % |
| Urban | 246 | 28 (11.4) | 218 (88.6) |
| Rural | 108 | 7 (6.5) | 101 (93.5) |
| Total | 354 | 35 (9.9) | 319 (90.1) |

P value=0.107



Dental caries

Regarding oral diseases, the study explores statistically significant relationship between dental caries and both oral protozoans, and the findings demonstrated that the higher positive rate of *E. gingivalis* (93.3%) was noticed among those with dental caries and it was encountered in 86.8% of subjects free from dental caries Table (3). While *T. tenax* infections were found to be positive in nineteen (11.5%) specimens from 165 patients with dental caries and in sixteen (8.5%) specimens from 189 healthy controls Table (4).

These findings are at variance with the studies of several investigators in the world like: in Kirkuk, [14] greatest percentage of both *E. gingivalis* and patients who had *T. tenax* with dental caries (29.62% and 25.92% respectively) but no infection was observed among those with no clinical signs, in Babylon, out of 59 specimens with dental caries, 22 cases were positive, and the highest rate of *T. tenax* was among the patients who suffered from dental caries (37.2%), but the rate of infection in the patient who have not shown any pathological conditions was (3%) in Al-Hilla city [15]. The [16] in Turkey, highlighted that dental caries was not found to be statistically significantly related with the presence of oral protozoa, [13] in Nigeria, observed 22.45% and 24.49% rates of infection with *E. gingivalis* and *T. tenax* respectively among patients with dental caries.

The differences in the research outcomes might be attributed to oral health issues.

Table (3) Occurrence of *Entamoeba gingivalis* among participants with Dental caries

| Oral disease | <i>Entamoeba gingivalis</i> | | |
|---------------|-----------------------------|---------------|--------------|
| | No. examined | No. +ve % | No. -ve % |
| Dental caries | 165 | 154 (93.3) | 11 (6.7) |
| Healthy mouth | 189 | 164 (86.8) | 25 (13.2) |
| Total | 354 | 318 (89.8) | 36 (10.2) |

Pvalue=0.03

Table (4) Occurrence of *Trichomonas tenax* among participants with dental caries

| Oral disease | <i>Trichomonas tenax</i> | | |
|---------------|--------------------------|--------------|---------------|
| | No. examined | No. +ve % | No. -ve % |
| Dental caries | 165 | 19 (11.5) | 146 (88.5) |
| Healthy mouth | 189 | 16 (8.5) | 173 (91.5) |
| Total | 354 | 35 (9.9) | 319 (90.1) |

P value=0.2171

Diabetes

Regarding diabetes, there is a statistically significant link between diabetes prevalence and *E. gingivalis* prevalence. Table (5) showed 97.1% and 35.7% infection rates of *E. gingivalis* among diabetics and non-diabetic participants respectively. This finding is disagreement with that of Chomicz et al. (2004) in Poland, who recorded that the frequency of parasitic protozoans, *T. tenax* and *E. gingivalis*, was higher in the control individual than in the diabetics (26.6% and 13.3%, respectively), and concluded it may be because metabolic illness that causes serious disturbances for multi-organ in addition to insulin therapy are the very important factors varying oral cavity ecology for people with diabetes and favor infections with other microorganisms. Diabetes is a significant metabolic condition that cause systemic problem that are also noticeable in the oral cavity [18]. While *T. tenax* was more common in individuals with non-diabetes (12.2%) compare to the diabetics (9.5%), but no relation was found as shown in Table (6).

There are contradicting reports in the literatures regarding diabetes: one study reports a high incidence of these commensals (74%) in adult diabetics [19] and another study reports a low incidence [20]. While other researchers stated that oral protozoa and diabetes were not shown to be significantly associated like: [21] in France, [22] in Turkey.

Table (5) Distribution of *Entamoeba gingivalis* between diabetics and non-diabetics

| | <i>Entamoeba gingivalis</i> | | |
|----------------------|-----------------------------|-----------------------|----------------------|
| | No. examined | No. +ve % | No. -ve % |
| Diabetics | 312 | 303 (97.1) | 9 2.8 |
| Non-diabetics | 42 | 15 (35.7) | 27 64.2 |
| Total | 354 | 318 (89.8) | 36 (10.2) |

P value=0.000

Table (6) Distribution of *T.tenax* between diabetics and non-diabetics

| | <i>T.tenax</i> | | |
|----------------------|----------------|---------------------|-----------------------|
| | No. examined | No. +ve % | No. -ve % |
| Diabetics | 305 | 29 (9.5) | 276 (90.4) |
| Non-diabetics | 49 | 6 (12.2) | 43 (87.7) |
| Total | 354 | 35 (9.9) | 319 (90.1) |

P value=0.351

Conflict of interests.

There are non-conflicts of interest.

References

- [1] Nadia Tawfiq Jaffer, Aleen Sardar Al-noori, Ahmed Mohammed Salih and Soleen Sardar Zuhdi. Molecular detection of *Trichomonas tenax* among individuals attending dental care units using PCR in Duhok city – Kurdistan region. Journal of University of Duhok, Vol. 22, No.2 (Pure and Eng. Sciences), Pp 195-202. 2019.
- [2] Ogunshe, A. and Odumesi, O. Users' perceptions and efficacy of indigenous adjunct teeth-cleansing agents on the bacterial flora of human dental caries. Afr. J. Clin. Exper. Microbiol., 11(3): 182-194. 2010.
- [3] Favoreto, J.S. and Machado, M.I. Incidence, morphology and diagnostic studies of *Entamoeba gingivalis*, Gros, 1849. Rev. Soc. Bras. Med. Trop., 28(4): 379-387. 1995.



- [4] Alaa Yaseen; Azmi Mahafzah; Deema Dababseh; Duaa Taim; Ahmad A. Hamdan; Esraa Al-Fraihat; Yazan Hassona; Gülşen Özkaya Şahin; Julien Santi-Rocca; and Malik Sallam. Oral Colonization by *Entamoeba gingivalis* and *Trichomonas tenax*: A PCR-Based Study in Health, Gingivitis, and Periodontitis. *Front. Cell. Infect. Microbiol.*, volume 11. 2021.
- [5] Sarowska, J.; Wonjnicz, D.; Kaczkowski, H. and Jankowski, S. The occurrence of *Entamoeba gingivalis* and *Trichomonas tenax* in patients with periodontal disease. *Adv Clin Exp Med*. J.13(2) :291-297. 2004.
- [6] Fahimeh Rashidi Maybodi, Ahmad Haerian Ardakani, Ali Fattahi Bafghi, Alireza Haerian Ardakani and Akram Zafarbakhsh. The Effect of Nonsurgical Periodontal Therapy on *Trichomonas Tenax* and *Entamoeba gingivalis* in Patients with Chronic Periodontitis. *J Dent Shiraz Univ Med Sci.*, 2016 September; 17(3): 171-176. 2016.
- [7] Ahmed, K.Al-hamiary, Mahdi Y. Kezar and Youniss A.AlKhafaji. Prevalence of Oral Protozoa in Periodontitis and Gingivitis Patients Who's Attended to Clinics Periodontics, Dentistry College\Babylon Univ. kufa unv. Magazen vol. (3), (1). 2011.
- [8] Zeibig, E.A. Clinical parasitology. A practical approach. Sanders an Imprint of Elsevier, Philadelphia 1997.
- [9] Roberts, L.S. and Janovy, J. Gerald, D. Schmidt and Larry, S. Roberts Foundations of parasitology, 8th Ed. McGraw – Hill companies, Inc. New York 2009.
- [10] Jia Yan, G.; QiuPing, L.; ZhiPing, C.; Hua, H.; Tao, J.; Wei, M. and XiFeng, T. Survey of *Entamoeba gingivalis* infection status in Tangshan. *Zhongguo Bingyuan Shengwuxue Zazhi / J. Patho. Biol.*, 4(4): 296-297, 2009.
- [11] Ozumba, U.C.; Ozumba, N. and Ndiokwelu, E.M. Oral protozoa in a Nigerea population. *Afr. J. Clin. Exper. Microbiol.*, 5(1): 15-19, 2004.
- [12] Mahdi, B.M.; Hazim, W.; Maki, B.; Mahdi, A.G.M. and Ameen, M.K. A Study on Humoral Immunity and Oral Bacterial Diversity in Patients with *Trichomonas tenax* Infection. *I. P. M. J.*, 8(4): 339-343, 2009.
- [13] Onyido, A.E.; Amadi, E.S.; Olofin, I.; Onwumma, A.A.; Okoh, I.C. and Chikwendu, C.I. Prevalence of *Entamoeba gingivalis* and *Trichomonas tenax* among dental patients attending Federal School of Dental Technology and Therapy clinic, Enugu, Nigeria. *Nature and Science*, 9(9): 59-62, 2011.
- [14] Kadir, M.A.; AL-Mashhadani, A.M.; Tahir, S.S. and Chapook, G.O. Prevalence of *Trichomonas tenax* and *Entamoeba gingivalis* among patients attending dental clinics in Kirkuk city. *Al-Taqani*, 20(1): 36-40, 2007.
- [15] Maaleh, S.F.A. Prevalence range of *Trichomonas tenax* among patients attended to dental clinics. *J. Babylon University/Pure and Applied Sciences*, 19(1): 97-99, 2011.
- [16] Ozcelik, S.; Gedik, T.; Gedik, R. and Malatyali, E. Investigation of the relationship between oral and dental health and presence of *Entamoeba gingivalis* and *Trichomonas tenax*. *Turkiye Parazitol. Derg.*, 34: 155-9, 2010.
- [17] Chomicz, L.; Szubińska, D.; Piekarczyk, J.; Wojtowicz, A.; Piekarczyk, B.; Starościak, B. and Fiedor, P. Occurrence of oral subclinical infections in insulin treated diabetics. *Wiad Parazytol.*, 50(2): 177-180, 2004.



- [18] Ziółkowska, J. Oral health status and dental service needs of diabetic patients. 52(3):103-14, 2006.
- [19] Nocito-Mendoza, I.; Vasconi-Correas, M.D.; Ponce de Leon-Horianski, P. and Zdero-Pandzich, M.. *Entamoeba gingivalis* and *Trichomonas tenax* in diabetic patients. RCOE., 8(1): 13-23, 2003.
- [20] Linke, H.; John, T. Gannon; J., and Jules, N.Obin. Clinical survey of *Entamoeba gingivalis* by multiple sampling in patients with advanced periodontal disease. International Journal for Parasitology. Volume 19, Issue 7, Pages 803-808, 1989.
- [21] Cambon, M.; Petavy, A.F.; Guitot, J.; Glandier, I.; Deguillaume, J. and Coulet, M.. A study of the frequency of protozoa and yeasts isolated from the paradontium of 509 subjects. Pathol. Biol., 27 (10): 603-606, 1979.
- [22] Ozcelik, S.; Gedik, T.; Gedik, R. and Malatyali, E. Investigation of the relationship between oral and dental health and presence of *Entamoeba gingivalis* and *Trichomonas tenax*. Turkiye Parazit. Derg., 34: 155-9, 2010.