

On Types of S_β - Functions

Hashmiya Ibrahim Nasser

Department of Math. - AL-Qadissiya Education

jaberabraham @ yahoo. com

Abstract

In this paper we introduce and study another types of functions in a topological spaces namely , S_β -compact and S_β -coercive by using the concept of S_β -open sets . Also we investigate some properties of these concepts and the relation between them.

Keywords: S_β -open, S_β -closed, S_β -compact space , S_β -continuous , S_β -compact, S_β -coercive.

الخلاصة

في هذا البحث قدمنا انواع من الدوال في الفضاءات التوبولوجية أسميناها الدوال (المرصوصة S_β -) و (الاضطرابية S_β -) باستخدام مفهوم المجموعة (S_β -open) وقدمنا بعض الخواص المبرهنات حول هذه الدوال والعلاقة بينهما. الكلمات المفتاحية: المفتوحة- S_β ، مغلقة - S_β ، الفضاء المضغوط- S_β ، المستمر- S_β ، المدمجة- S_β ، القسرية- S_β .

Introduction

In 1963, Levine initiated the notation of semi-open (briefly S -open) sets and study their properties of this concept. Throughout the properties in topological spaces. In 1982 , AbdEl-monsef defined the class of β -open set. Also in 2012, Kalaf B. and others were introduced a new concept denoted by S_β -open set. Finally, in 2013 , Kalaf B. and Ahmed K. wre introduced a new type of compact spaces namely S_β -compact. Present paper, (X, T) and (Y, T') (or simply X and Y) denote topological spaces . The closure (resp. interior) of a subset A of a space X which denoted by $cl(A)$ (resp. $int(A)$) . A subset A of X is called S -open (resp. β -open) , if $A \subseteq cl(int(A))$ (resp. $A \subseteq cl(int(cl(A)))$) . The complement of S -open (resp. β -open) set is called S -closed (resp. β -closed) set.

Finally in section two , we give some basic properties of new types of functions and relation between them .

1. Basic definitions and notations:

We introduce some elementary concept which we need in our work.

1.1. Definition: [Al-Sheikhly, 2003]

A topological space X is called:

- i. Locally indiscrete if every open subset of X is closed.
- ii. Hyper-connected if every non-empty open subset of is dense.

1.2. Definition: [Kalaf, 2012]

- i. A S -open subset A of a topological space X is called S_β -open if for all $x \in A$, there is a β -closed set F such that $x \in F \subseteq A$. A complement of S_β -open is called S_β -closed.
- ii. A subset A of a topological space X is called S_β -open if and only if A is S -open and it is a union of β -closed sets.
- iii. A subset N of a topological space X is called S_β -neighborhood of a subset A of X , if there is a S_β -open set U such that $A \in U \subseteq N$. When $A = \{x\}$ we say that N is S_β - neighborhood of x .

iv. A topological space X is called \mathcal{S}_β -compact, if for every \mathcal{S}_β -open cover of X has finite subcover. Clearly, every \mathcal{S} -compact space is \mathcal{S}_β -compact.

1.3. Remark: [Kalaf, 2013]

- i. If X is a T_1 -space, then every \mathcal{S}_β -open subset in X is \mathcal{S} -open.
- ii. A space X is hyper-connected if and only if a \mathcal{S}_β -open subsets of X are \emptyset and X .
- iii. If a space X is locally indiscrete, then every \mathcal{S} -open subset of X is \mathcal{S}_β -open.

1.4. Theorem: [Kalaf, 2012]

- i. If B is clopen subset of a topological space X and A is open, then $A \cap B$ is \mathcal{S}_β -open.
- ii. Let $A \subseteq Y \subseteq X$, if A is \mathcal{S}_β -open subset in X and Y is open subset in X , then A is \mathcal{S}_β -open subset in Y .
- iii. Let $A \subseteq Y \subseteq X$, if A is \mathcal{S}_β -open subset in Y and Y is clopen subset in X , then A is \mathcal{S}_β -open subset in X .

1.5. Definition: [Al-Sheikhly, 2003; Kalaf, 2013]

Let $f: X \rightarrow Y$ be a function, then f is called:

- i. \mathcal{S} -compact if the inverse image for every compact set in Y is \mathcal{S} -compact set in X .
- ii. \mathcal{S}_β -closed if the image for every closed set in Y is \mathcal{S}_β -closed set in X .
- iii. \mathcal{S}_β -irresolute if the inverse image for every \mathcal{S}_β -open set in Y is \mathcal{S}_β -open set in X .

2. Type of \mathcal{S}_β -Functions:

In this section, we introduce a new \mathcal{S}_β -functions called \mathcal{S}_β -compact and \mathcal{S}_β -coercive functions.

2.1. Definition:

A function $f: X \rightarrow Y$ is said to be \mathcal{S}_β -compact if the inverse image for every compact set in Y is \mathcal{S}_β -compact set in X .

2.2. Example:

- i. An identity function $f: (X, T) \rightarrow (X, T')$ with $X = \mathcal{R}$, $T = \{\emptyset, X, \{0\}\}$ and $T' = T_{ind}$ is \mathcal{S}_β -compact.
- ii. Let $X = \{1, 2, 3\}$ and $Y = \{2, 4, 6\}$ with topologies $T = T_{ind}$, $T' = \{\emptyset, Y, \{4\}\}$ resp. A function $f: X \rightarrow Y$ defined by $f(x) = 2x, \forall x \in X$ is \mathcal{S}_β -compact, since X is locally indiscrete space [by using remark (1.3.iii)].

The following example shows that not every function is \mathcal{S}_β -compact.

2.3. Example:

Consider a countable set X with co-countable topology, then the constant function from X into any space Y is not \mathcal{S}_β -compact.

2.4. Theorem:

Every \mathcal{S} -compact function is \mathcal{S}_β -compact.

Proof: By using definition (1.2.iv), this is just the condition of our theorem.

The converse of the Theorem above is not true in general as the example shows:

2.5. Example:

Let $X = \mathcal{R}$ with topology $T = \{\emptyset, X, \{0\}\}$, then a function $f: (X, T) \rightarrow (X, T_U)$ which defined by $f(x) = 0, \forall x \in X$ is \mathcal{S}_β -compact, but not \mathcal{S} -compact.

In general a \mathcal{S}_β -compact and compact functions are independent as the following examples:

2.6. Example:

A function in example (2.3) is compact, but not \mathcal{S}_β -compact.

2.7. Example:

Let $X = (0,1)$ with topology $\tau = \{\emptyset, X, G = (0, 1 - \frac{1}{n})\}, n = 2, 3, \dots$. A constant function f from a space X is a S_β -compact, but not its compact.

Recall that every S -compact function is compact [Al-Sheikhly H. 2003], the converse is not true as the following example shows:

2.8. Example:

Let $X = I \cup \{x\}$ with I uncountable set and $x \notin I$, let $\tau = \{\emptyset, X, \{x\}\}$ be a topology on X , a constant function from X into itself is compact but not S -compact, because X is not S -compact space, since $\{\{x, x_i\} : i \in I\}$ is a S -covering of X but has no finite subcover.

2.9. Theorem:

Let $f: X \rightarrow Y$ be a S_β -compact function and A be a clopen subset of X , then $f|_A: A \rightarrow Y$ is also S_β -compact.

Proof:

Let K be a compact subset of Y , then $f^{-1}(K)$ is S_β -compact set in X , since A clopen in X . Then by Theorem (1.4.iii), $A \cap f^{-1}(K)$ is a S_β -compact set in A , but $A \cap f^{-1}(K) = f|_A^{-1}(K)$, then $f|_A$ is S_β -compact.

2.10. Remark:

A composition of two S_β -compact functions not necessary S_β -compact.

2.11. Theorem:

Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be two functions, then:

- i. If f and g are S_β -compact and Y be a locally indiscrete, then $g \circ f$ is S_β -compact.
- ii. If f and g are S_β -compact and Y be a T_1 -space, then $g \circ f$ is S_β -compact.
 - iii. If g compact and f be a S_β -compact, then $g \circ f$ is S_β -compact.
 - iv. If g be a S -compact and f be a S_β -compact, then $g \circ f$ is S_β -compact.

2.12. Definition:

A function $f: X \rightarrow Y$ is said to be S_β -coercive if for every S_β -compact subset B of Y there is S_β -compact subset A of X such that $f(X/A) \subseteq (Y/B)$.

2.13. Example:

- i. The identity function for any space is S_β -coercive.
- ii. Let $X = \{1, 2, 3\}, Y = \{4, 5\}, T_X = \{\emptyset, X, \{3\}\}, T_Y = T_{ind}$ and $f: X \rightarrow Y$ be a function which defined by $f(1) = f(2) = 4, f(3) = 5$, then f is S_β -coercive.

2.14. Theorem:

A S_β -compact function from S_β -compact space is S_β -coercive.

Proof:

Let $f: X \rightarrow Y$ be a S_β -compact function and let B be a S_β -compact subset of Y . Since X be a S_β -compact. Then $f(X/X) = \emptyset \subseteq f(Y/B)$, thus f is S_β -coercive.

2.15. Theorem:

A restriction S_β -coercive function on clopen subset is S_β -coercive.

Proof:

Let $f: X \rightarrow Y$ be a S_β -coercive function and F be a clopen subset of X , to show that $f|_F: F \rightarrow Y$ is a S_β -coercive function, let B be a S_β -compact subset of Y . Then there is a S_β -compact subset A of X such that $f(X/A) \subseteq (Y/B)$. Since F be a clopen subset of X , then by Theorem (1.4.iii), $F \cap A$ is S_β -compact subset of F . Since $f|_F(F \cap A) = f(F/A)$ and $F/A \subseteq X/A$
 $\Rightarrow f(F/A) \subseteq f(X/A) \Rightarrow f|_F(F/F \cap A) \subseteq Y/B$, hence $f|_F: F \rightarrow Y$ is S_β -coercive function.

2.16. Theorem:

A composition of two \mathcal{S}_β -coercive functions is \mathcal{S}_β -coercive.

Proof: Clear.

Recall that the \mathcal{S}_β -irresolute image of \mathcal{S}_β -compact is \mathcal{S}_β -compact [Kalaf K. 2012].

2.17. Theorem:

Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be two functions such that:

- i. If $g \circ f$ is \mathcal{S}_β -coercive with g is \mathcal{S}_β -irresolute and one to one, then f is \mathcal{S}_β -coercive .
- ii. If $g \circ f$ is \mathcal{S}_β -coercive with f is \mathcal{S}_β -irresolute and onto, then g is \mathcal{S}_β -coercive.

Proof:

i. Let B be a \mathcal{S}_β -compact subset of Y , then $g(B)$ is a \mathcal{S}_β -compact subset of Z . Since $g \circ f$ be a \mathcal{S}_β -coercive , then there is a \mathcal{S}_β -compact subset A of X such that $g \circ f (X/A) \subseteq Z / g(B)$, then:

$$f(X/A) = g^{-1}(g \circ f (X/A)) \subseteq g^{-1}(Z/g(B)) = g^{-1}(Z \cap (g(B))^c) = g^{-1}(Z) \cap g^{-1}(g(B)^c) = Y/B$$

,thus

f is \mathcal{S}_β -coercive function.

ii. Let C be a \mathcal{S}_β -compact subset of Z . Since $g \circ f$ is \mathcal{S}_β -coercive , then there is a \mathcal{S}_β -compact subset A of X such that $g \circ f (X/A) \subseteq Z / C$, thus $g(f(A^c)) \subseteq Z / C$, since f onto we get $g((f(A))^c) \subseteq Z / C$. Then $f(A)$ is \mathcal{S}_β -compact subset of Y . Thus g is \mathcal{S}_β -coercive function.

Reference:

Al-Sheikhly A. H., 2003, " On Semi Proper G-Space ", Thesis submitted to the col. of science AL-Mustansirya univ.

Kalaf B. A. and N. K. Ahmed, 2012," \mathcal{S}_β -open and \mathcal{S}_β -continuity in topological spaces " Int. J. of Math. Sci. and Engg. Appls. , III(1), P. 1-13.

Kalaf B. A. and N. K. Ahmed, 2012, " \mathcal{S}_β -compact and \mathcal{S}_β -compact spaces " Int. J. of Math. Sci. and Engg. Appls. , Vol. 3 , Issue 4 , April.

Kalaf B. A. and N. K. Ahmed ,2012, " Weak separation Axioms And Functions With \mathcal{S}_β -closed Graph " , Int. J. of Math. Sci. and Engg. Appls. , Vol. 6 , No. III .

Kalaf B. A. and N. K. Ahmed, 2013, " \mathcal{S}_β -paracompact Spaces ", J. of Ad. Stud. in topology , Vol. 4 , No. 1 ,p. 40-47.

Kalaf B. A. and N. K. Ahmed, 2013," \mathcal{S}_β -compact Sets and \mathcal{S}_β -locally Compact Spaces ", J. of Ad. Stud. in topology, Vol. 4 , No. 1, p. 113-118.

Levine N., 1963, " Semi open sets and semi contiuous in topological spaces " , Amer. Math. , Monthly 70 , 36-41.

Mashhour A. S. , Abd El-monsef M. E. and El-Deeb S. E., 1982, " On pre-continuous and week Pre-continuous mappings", proc. Math. Phys. Soc., Egypt 53, p.47-53.

